

Top-Antitop Production at Hadron Colliders

Roberto BONCIANI

*Laboratoire de Physique Subatomique et de Cosmologie,
Université Joseph Fourier/CNRS-IN2P3/INPG,
F-38026 Grenoble, France*



Plan of the Talk

- General Introduction
 - Top Quark at the Tevatron
 - LHC Perspectives
- Status of the Theoretical calculations
 - The General Framework
 - Total Cross Section at NLO
- Analytic Two-Loop QCD Corrections
- Conclusions

Top Quark


Top Quark

- With a mass of $m_t = 173.1 \pm 1.3 \text{ GeV}$, the TOP quark (the up-type quark of the third generation) is the heaviest elementary particle produced so far at colliders.
- Because of its mass, top quark is going to play a unique role in understanding the EW symmetry breaking $\Rightarrow$ **Heavy-Quark physics crucial at the LHC.**

- Two production mechanisms

- $pp(\bar{p}) \rightarrow t\bar{t}$


- $pp(\bar{p}) \rightarrow t\bar{b}, tq'(\bar{q}'), tW^-$



- Top quark does not hadronize, since it decays in about $5 \cdot 10^{-25} \text{ s}$ (one order of magnitude smaller than the hadronization time) $\Rightarrow$ opportunity to study the quark as single particle

- Spin properties
- Interaction vertices
- Top quark mass

- Decay products: almost exclusively $t \rightarrow W^+ b$ ($|V_{tb}| \gg |V_{td}|, |V_{ts}|$)




Top Quark

- With a mass of $m_t = 173.1 \pm 1.3 \text{ GeV}$, the TOP quark (the up-type quark of the third generation) is the heaviest elementary particle produced so far at colliders.
- Because of its mass, top quark is going to play a unique role in understanding the EW symmetry breaking $\Rightarrow$ **Heavy-Quark physics crucial at the LHC.**


Two production mechanisms

- $pp(\bar{p}) \rightarrow t\bar{t}$



Pair Production

- $pp(\bar{p}) \rightarrow t\bar{b}, tq'(\bar{q}'), tW^-$




Single Top

- Top quark does not hadronize, since it decays in about $5 \cdot 10^{-25} \text{ s}$ (one order of magnitude smaller than the hadronization time) $\Rightarrow$ opportunity to study the quark as single particle

- Spin properties
- Interaction vertices
- Top quark mass

- Decay products: almost exclusively $t \rightarrow W^+ b$ ($|V_{tb}| \gg |V_{td}|, |V_{ts}|$)




Top Quark

- With a mass of $m_t = 173.1 \pm 1.3 \text{ GeV}$, the TOP quark (the up-type quark of the third generation) is the heaviest elementary particle produced so far at colliders.
- Because of its mass, top quark is going to play a unique role in understanding the EW symmetry breaking $\Rightarrow$ **Heavy-Quark physics crucial at the LHC.**

- Two production mechanisms

- $pp(\bar{p}) \rightarrow t\bar{t}$


- $pp(\bar{p}) \rightarrow t\bar{b}, tq'(\bar{q}'), tW^-$



- Top quark does not hadronize, since it decays in about $5 \cdot 10^{-25} \text{ s}$ (one order of magnitude smaller than the hadronization time) $\Rightarrow$ opportunity to study the quark as single particle

- Spin properties
- Interaction vertices
- Top quark mass

- Decay products: almost exclusively $t \rightarrow W^+ b$ ($|V_{tb}| \gg |V_{td}|, |V_{ts}|$)



Top Quark


Tevatron

- To date the Top quark could be produced and studied only at the Tevatron (discovery 1995)
- $p\bar{p}$ collisions at $\sqrt{s} = 1.96$ TeV
- $L \sim 6.5\text{fb}^{-1}$ reached in 2009
- $\mathcal{O}(10^3)$ $t\bar{t}$ pairs produced so far
- Only recently confirmation of single-t



LHC

- Running since end 2009
- pp collisions at $\sqrt{s} = 7$ (14) TeV
- LHC will be a factory for heavy quarks ($\mathcal{L} \sim 10^{33}-10^{34}\text{cm}^{-2}\text{s}^{-1}$, $t\bar{t}$ at $\sim 1\text{Hz}$!)
- Even in the first low-luminosity phase (2 years $\sim 1\text{fb}^{-1}$ @ 7 TeV) $\sim \mathcal{O}(10^4)$ registered $t\bar{t}$ pairs



Top Quark @ Tevatron

Top Quark @ Tevatron

Events measured at Tevatron

$$\sigma_{t\bar{t}} \sim 7\text{pb}$$

$$\left[\begin{array}{l} \bullet \quad p\bar{p} \rightarrow t\bar{t} \rightarrow W^+ b W^- \bar{b} \rightarrow l\nu l\nu b\bar{b} \\ \bullet \quad p\bar{p} \rightarrow t\bar{t} \rightarrow W^+ b W^- \bar{b} \rightarrow l\nu q\bar{q}' b\bar{b} \\ \bullet \quad p\bar{p} \rightarrow t\bar{t} \rightarrow W^+ b W^- \bar{b} \rightarrow q\bar{q}' q\bar{q}' b\bar{b} \end{array} \right]$$

Dilepton $\sim 10\%$


Lep+jets $\sim 44\%$

All jets $\sim 46\%$

Top Quark @ Tevatron

Events measured at Tevatron

$$\sigma_{t\bar{t}} \sim 7\text{pb}$$


$$p\bar{p} \rightarrow t\bar{t} \rightarrow W^+bW^-\bar{b} \rightarrow l\nu l\nu b\bar{b}$$

$$p\bar{p} \rightarrow t\bar{t} \rightarrow W^+bW^-\bar{b} \rightarrow l\nu q\bar{q}'b\bar{b}$$

$$p\bar{p} \rightarrow t\bar{t} \rightarrow W^+bW^-\bar{b} \rightarrow q\bar{q}'q\bar{q}'b\bar{b}$$

Dilepton $\sim 10\%$

Lep+jets $\sim 44\%$


All jets $\sim 46\%$

2 high- p_T lept, ≥ 2 jets and ME

Top Quark @ Tevatron

Events measured at Tevatron

$$\sigma_{t\bar{t}} \sim 7\text{pb}$$



- $p\bar{p} \rightarrow t\bar{t} \rightarrow W^+bW^-\bar{b} \rightarrow l\nu l\nu b\bar{b}$
- $p\bar{p} \rightarrow t\bar{t} \rightarrow W^+bW^-\bar{b} \rightarrow l\nu q\bar{q}'b\bar{b}$
- $p\bar{p} \rightarrow t\bar{t} \rightarrow W^+bW^-\bar{b} \rightarrow q\bar{q}'q\bar{q}'b\bar{b}$

Dilepton $\sim 10\%$

Lep+jets $\sim 44\%$

All jets $\sim 46\%$

2 high- p_T lept, ≥ 2 jets and ME

1 isol high- p_T lept, ≥ 4 jets and ME

Top Quark @ Tevatron

Events measured at Tevatron

$$\sigma_{t\bar{t}} \sim 7\text{pb}$$


$$p\bar{p} \rightarrow t\bar{t} \rightarrow W^+bW^-\bar{b} \rightarrow l\nu l\nu b\bar{b}$$

Dilepton $\sim 10\%$


$$p\bar{p} \rightarrow t\bar{t} \rightarrow W^+bW^-\bar{b} \rightarrow l\nu q\bar{q}'b\bar{b}$$

Lep+jets $\sim 44\%$


$$p\bar{p} \rightarrow t\bar{t} \rightarrow W^+bW^-\bar{b} \rightarrow q\bar{q}'q\bar{q}'b\bar{b}$$

All jets $\sim 46\%$

2 high- p_T lept, ≥ 2 jets and ME

NO lept, ≥ 6 jets and low ME

1 isol high- p_T lept, ≥ 4 jets and ME

Top Quark @ Tevatron

Events measured at Tevatron

$$\sigma_{t\bar{t}} \sim 7\text{pb}$$

$$\left[\begin{array}{l} \bullet \quad p\bar{p} \rightarrow t\bar{t} \rightarrow W^+ b W^- \bar{b} \rightarrow l\nu l\nu b\bar{b} \\ \bullet \quad p\bar{p} \rightarrow t\bar{t} \rightarrow W^+ b W^- \bar{b} \rightarrow l\nu q\bar{q}' b\bar{b} \\ \bullet \quad p\bar{p} \rightarrow t\bar{t} \rightarrow W^+ b W^- \bar{b} \rightarrow q\bar{q}' q\bar{q}' b\bar{b} \end{array} \right.$$

Dilepton $\sim 10\%$

Lep+jets $\sim 44\%$


All jets $\sim 46\%$

2 high- p_T lept, ≥ 2 jets and ME


NO lept, ≥ 6 jets and low ME

1 isol high- p_T lept, ≥ 4 jets and ME


Background Processes




W+jets



QCD



Drell-Yan



Di-boson

Top Quark @ Tevatron

Events measured at Tevatron

$$\sigma_{t\bar{t}} \sim 7\text{pb}$$

$$\left[\begin{array}{l} \bullet \quad p\bar{p} \rightarrow t\bar{t} \rightarrow W^+ b W^- \bar{b} \rightarrow l\nu l\nu b\bar{b} \\ \bullet \quad p\bar{p} \rightarrow t\bar{t} \rightarrow W^+ b W^- \bar{b} \rightarrow l\nu q\bar{q}' b\bar{b} \\ \bullet \quad p\bar{p} \rightarrow t\bar{t} \rightarrow W^+ b W^- \bar{b} \rightarrow q\bar{q}' q\bar{q}' b\bar{b} \end{array} \right.$$

Dilepton $\sim 10\%$

Lep+jets $\sim 44\%$


All jets $\sim 46\%$

2 high- p_T lept, ≥ 2 jets and ME

NO lept, ≥ 6 jets and low ME

1 isol high- p_T lept, ≥ 4 jets and ME

Background Processes




Top Quark @ Tevatron

● Total Cross Section

$$\sigma_{t\bar{t}} = \frac{N_{data} - N_{bkg}}{\epsilon L}$$

Combination CDF-D0 ($m_t = 175 \text{ GeV}$)

$$\sigma_{t\bar{t}} = 7.0 \pm 0.6 \text{ pb} \quad (\Delta\sigma_{t\bar{t}}/\sigma_{t\bar{t}} \sim 9\%)$$




● Top-quark Mass

- Fundamental parameter of the SM. A precise measurement useful to constraint Higgs mass from radiative corrections (Δr)
- A possible extraction: $\sigma_{t\bar{t}} \Rightarrow$ need of precise theoretical determination

$$\frac{\Delta m_t}{m_t} \sim \frac{1}{5} \frac{\Delta\sigma_{t\bar{t}}}{\sigma_{t\bar{t}}}$$

Combination CDF-D0

$$m_t = 173.1 \pm 1.3 \text{ GeV} (0.75\%)$$




Top Quark @ Tevatron

- W helicity fractions $F_i = B(t \rightarrow bW^+(\lambda_W = i))$ ($i = -1, 0, 1$) measured fitting the distribution in θ^* (the angle between l^+ in the W^+ rest frame and W^+ direction in the t rest frame)

$$\frac{1}{\Gamma} \frac{d\Gamma}{d\cos\theta^*} = \frac{3}{4}F_0 \sin^2\theta^* + \frac{3}{8}F_-(1 - \cos\theta^*)^2 + \frac{3}{8}F_+(1 + \cos\theta^*)^2$$

$$F_0 + F_+ + F_- = 1$$

$$F_0 = 0.66 \pm 0.16 \pm 0.05 \quad F_+ = -0.03 \pm 0.06 \pm 0.03$$



- Spin correlations measured fitting the double distribution (θ_1 (θ_2) is the angle between the dir of flight of l_1 (l_2) in the $t(\bar{t})$ rest frame and the $t(\bar{t})$ direction in the $t\bar{t}$ rest frame)

$$\frac{1}{N} \frac{d^2 N}{d\cos\theta_1 d\cos\theta_2} = \frac{1}{4}(1 + \kappa \cos\theta_1 \cos\theta_2)$$

$$-0.455 < \kappa < 0.865 \text{ (68\% CL)}$$

- Forward-Backward Asymmetry

$$A_{FB} = \frac{N(y_t > 0) - N(y_t < 0)}{N(y_t > 0) + N(y_t < 0)}$$

$$A_{FB} = (19.3 \pm 6.5(\text{sta}) \pm 2.4(\text{sys}))\%$$

Top Quark @ Tevatron

Tevatron searches of physics BSM in top events

- New production mechanisms via new spin-1 or spin-2 resonances: $q\bar{q} \rightarrow Z' \rightarrow t\bar{t}$ in lepton+jets and all hadronic events (bumps in the invariant-mass distribution)

- Top charge measurements (recently excluded $Q_t = -4/3$)

- Anomalous couplings

$$L = -\frac{g}{\sqrt{2}} \bar{b} \left\{ \gamma^\mu (V_L P_L + V_R P_R) + \frac{i\sigma^{\mu\nu} (p_t - p_b)_\nu}{M_W} (g_L P_L + g_R P_R) \right\} t W_\mu^-$$

- From helicity fractions

- From asymmetries in the final state (for instance $A_{FB} = 3/4 (F_+ - F_-)$)

- Forward-backward asymmetry

- Non SM Top decays. Search for charged Higgs: $t \rightarrow H^+ b \rightarrow q\bar{q}' b (\tau \nu b)$

- Search for heavy $t' \rightarrow W^+ b$ in lepton+jets

Top Quark @ Tevatron

Tevatron searches of physics BSM in top events

- New physics in lepton+jet

- Top charge asymmetry

- Anomalous top production

$L = -$

- Forward

- Forward

- Forward

- Non SM Top decays. Search for charged Higgs: $t \rightarrow H^+ b \rightarrow q\bar{q}'b(\tau\nu b)$

- Search for heavy $t' \rightarrow W^+ b$ in lepton+jets

No Evidence
of New Physics so far

$\rightarrow t\bar{t}$ in

))

Top Quark @ Tevatron

	Value	Lum fb^{-1}	SM value	SM-like?
m_t	$173.1 \pm 0.6 \pm 1.1$ GeV	up to 4.8	/	/
$\sigma_{t\bar{t}}$	$7.0 \pm 0.3 \pm 0.4 \pm 0.4$ pb ($m_t = 175$ GeV)	2.8	6.7 pb	YES
W-helicity	$F^0 = 0.66 \pm 0.16 \pm 0.05$ $F^+ = -0.03 \pm 0.06 \pm 0.03$	1.9	$F^0 = 0.7$ $F^+ = 0$	YES
Spin Correlat.	$-0.455 < \kappa < 0.865$ (68% CL)	2.8	$\kappa = 0.8$	YES
A_{FB}	$0.19 \pm 0.07 \pm 0.02$	3.2	0.05 @NLO	YES
Γ_t	< 13.1 GeV (95% CL)	1.0	1.5 GeV	YES
τ_t	$c\tau_t < 52.5$ μ m (95% CL)	0.3	$\sim 10^{-16}$ m	YES
BR	$(t \rightarrow Wb)/(t \rightarrow Wq) > 0.61$ (95% CL)	0.2	$\sim 100\%$	YES
Charge	Exclude $Q_t = -4/3$ (87% CL)	1.5	2/3	YES

LHC Perspectives

● Cross Section

- With 100 pb^{-1} of accumulated data an error of $\Delta\sigma_{t\bar{t}}/\sigma_{t\bar{t}} \sim 15\%$ is expected (dominated by statistics!)
- After 5 years of data taking an error of $\Delta\sigma_{t\bar{t}}/\sigma_{t\bar{t}} \sim 5\%$ is expected

● Top Mass

- With 1 fb^{-1} Mass accuracy: $\Delta m_t \sim 1 - 3 \text{ GeV}$

● Top Properties

- W helicity fractions and spin correlations with $10 \text{ fb}^{-1} \Rightarrow 1\text{-}5\%$
- Top-quark charge. With 1 fb^{-1} we could be able to determine $Q_t = 2/3$ with an accuracy of $\sim 15\%$

● Sensitivity to new physics

- all the above mentioned points
- Narrow resonances: with 1 fb^{-1} possible discovery of a Z' of $M_{Z'} \sim 700 \text{ GeV}$ with $\sigma_{pp \rightarrow Z' \rightarrow t\bar{t}} \sim 11 \text{ pb}$

LHC Perspectives

● Cross Section

- With 100 pb^{-1} of accumulated data an error of $\Delta\sigma_{t\bar{t}}/\sigma_{t\bar{t}} \sim 15\%$ is expected (dominated by statistics!)
- After 5 years of data taking an error of $\Delta\sigma_{t\bar{t}}/\sigma_{t\bar{t}} \sim 5\%$ is expected

● Top Mass

- With 1 fb^{-1} Mass accuracy: $\Delta m_t \sim 1 - 3 \text{ GeV}$

● Top Properties

- W helicity fractions and spin correlations with $10 \text{ fb}^{-1} \Rightarrow 1\text{-}5\%$
- Top-quark charge. With 1 fb^{-1} we could be able to determine $Q_t = 2/3$ with an accuracy of $\sim 15\%$


● Sensitivity to new physics

- all the above mentioned points
- Narrow resonances: with 1 fb^{-1} possible discovery of a Z' of $M_{Z'} \sim 700 \text{ GeV}$ with $\sigma_{pp \rightarrow Z' \rightarrow t\bar{t}} \sim 11 \text{ pb}$

Top-Anti Top Pair Production

Top-Anti Top Pair Production

According to the factorization theorem, the process $h_1 + h_2 \rightarrow t\bar{t} + X$ can be sketched as in the figure:




$$\sigma_{h_1, h_2}^{t\bar{t}} = \sum_{i,j} \int_0^1 dx_1 \int_0^1 dx_2 f_{h_1,i}(x_1, \mu_F) f_{h_2,j}(x_2, \mu_F) \hat{\sigma}_{ij}(\hat{s}, m_t, \alpha_s(\mu_R), \mu_F, \mu_R)$$

$$s = (p_{h_1} + p_{h_2})^2, \quad \hat{s} = x_1 x_2 s$$


Top-Anti Top Pair Production

According to the factorization theorem, the process $h_1 + h_2 \rightarrow t\bar{t} + X$ can be sketched as in the figure:



Top-Anti Top Pair Production

According to the factorization theorem, the process $h_1 + h_2 \rightarrow t\bar{t} + X$ can be sketched as in the figure:




$$\sigma_{h_1, h_2}^{t\bar{t}} = \sum_{i,j} \int_0^1 dx_1 \int_0^1 dx_2 f_{h_1,i}(x_1, \mu_F) f_{h_2,j}(x_2, \mu_F) \hat{\sigma}_{ij}(\hat{s}, m_t, \alpha_s(\mu_R), \mu_F, \mu_R)$$

$$s = (p_{h_1} + p_{h_2})^2, \quad \hat{s} = x_1 x_2 s$$

The Cross Section: LO

The Cross Section: LO


$$q(p_1) + \bar{q}(p_2) \longrightarrow t(p_3) + \bar{t}(p_4)$$



Dominant at Tevatron

$\sim 85\%$

$$g(p_1) + g(p_2) \longrightarrow t(p_3) + \bar{t}(p_4)$$



Dominant at LHC

$\sim 90\%$

$$\sigma_{t\bar{t}}^{LO}(LHC, m_t = 171 \text{ GeV}) = 583 \text{ pb} \pm 30\%$$

$$\sigma_{t\bar{t}}^{LO}(Tev, m_t = 171 \text{ GeV}) = 5.92 \text{ pb} \pm 44\%$$

The Cross Section: NLO

The Cross Section: NLO

Fixed Order

- The NLO QCD corrections are quite sizable: + 25% at Tevatron and +50% at LHC.
Scale variation $\pm 15\%$.

Nason, Dawson, Ellis '88-'90; Beenakker, Kuijf, van Neerven, Smith '89-'91;
Mangano, Nason, Ridolfi '92; Frixione et al. '95; Czakon and Mitov '08
Melnikov and Schulze '09; Bernreuther and Si '10

- Mixed NLO QCD-EW corrections are small: - 1% at Tevatron and -0.5% at LHC.

Beenakker *et al.* '94 Bernreuther, Fuecker, and Si '05-'08
Kühn, Scharf, and Uwer '05-'06; Moretti, Nolten, and Ross '06.

The Cross Section: NLO

Fixed Order

- The NLO QCD corrections are quite sizable: + 25% at Tevatron and +50% at LHC. Scale variation $\pm 15\%$.

Nason, Dawson, Ellis '88-'90; Beenakker, Kuijf, van Neerven, Smith '89-'91; Mangano, Nason, Ridolfi '92; Frixione et al. '95; Czakon and Mitov '08
Melnikov and Schulze '09; Bernreuther and Si '10

- Mixed NLO QCD-EW corrections are small: - 1% at Tevatron and -0.5% at LHC.

Beenakker *et al.* '94 Bernreuther, Fuecker, and Si '05-'08
Kühn, Scharf, and Uwer '05-'06; Moretti, Nolten, and Ross '06.

- The QCD corrections to processes involving at least two large energy scales ($\hat{s}, m_t^2 \gg \Lambda_{QCD}^2$) are characterized by a logarithmic behavior in the vicinity of the boundary of the phase space

$$\sigma \sim \sum_{n,m} C_{n,m} \alpha_S^n \ln^m(1 - \rho) \quad m \leq 2n$$

The Cross Section: NLO

Fixed Order

- The NLO QCD corrections are quite sizable: + 25% at Tevatron and +50% at LHC. Scale variation $\pm 15\%$.

Nason, Dawson, Ellis '88-'90; Beenakker, Kuijf, van Neerven, Smith '89-'91;
Mangano, Nason, Ridolfi '92; Frixione et al. '95; Czakon and Mitov '08
Melnikov and Schulze '09; Bernreuther and Si '10

- Mixed NLO QCD-EW corrections are small: - 1%

Beenakker *et al.* '94 Bernreuther
Kühn, Scharf, and Uwer '05-'06

Inelasticity parameter

$$\rho = \frac{4m_t^2}{\hat{s}} \rightarrow 1$$

- The QCD corrections to processes involving at least two large energy scales ($\hat{s}, m_t^2 \gg \Lambda_{QCD}^2$) are characterized by a logarithmic behavior in the vicinity of the boundary of the phase space

$$\sigma \sim \sum_{n,m} C_{n,m} \alpha_S^n \ln^m (1 - \rho) \quad m \leq 2n$$

The Cross Section: NLO

Fixed Order

- The NLO QCD corrections are quite sizable: + 25% at Tevatron and +50% at LHC. Scale variation $\pm 15\%$.

Nason, Dawson, Ellis '88-'90; Beenakker, Kuijf, van Neerven, Smith '89-'91;
Mangano, Nason, Ridolfi '92; Frixione et al. '95; Czakon and Mitov '08
Melnikov and Schulze '09; Bernreuther and Si '10

- Mixed NLO QCD-EW corrections are small: - 1%

Beenakker *et al.* '94 Bernreuther
Kühn, Scharf, and Uwer '05-'06

Inelasticity parameter

$$\rho = \frac{4m_t^2}{\hat{s}} \rightarrow 1$$

- The QCD corrections to processes involving at least two large energy scales ($\hat{s}, m_t^2 \gg \Lambda_{QCD}^2$) are characterized by a logarithmic behavior in the vicinity of the boundary of the phase space

$$\sigma \sim \sum_{n,m} C_{n,m} \alpha_S^n \ln^m (1 - \rho) \quad m \leq 2n$$

- Even if $\alpha_S \ll 1$ (perturbative region) we can have at all orders

$$\alpha_S^n \ln^m (1 - \rho) \sim \mathcal{O}(1)$$

Resummation $\Rightarrow$ improved perturbation theory

The Cross Section: NLO

Fixed Order

- The NLO QCD corrections are quite sizable: + 25% at Tevatron and +50% at LHC. Scale variation $\pm 15\%$.

Nason, Dawson, Ellis '88-'90; Beenakker, Kuijf, van Neerven, Smith '89-'91; Mangano, Nason, Ridolfi '92; Frixione et al. '95; Czakon and Mitov '08
Melnikov and Schulze '09; Bernreuther and Si '10

- Mixed NLO QCD-EW corrections are small: - 1% at Tevatron and -0.5% at LHC.

Beenakker *et al.* '94 Bernreuther, Fuecker, and Si '05-'08
Kühn, Scharf, and Uwer '05-'06; Moretti, Nolten, and Ross '06.

All-order Soft-Gluon Resummation

- Leading-Logs (LL)

Laenen et al. '92-'95; Berger and Contopanagos '95-'96; Catani et al. '96.

- Next-to-Leading-Logs (NLL)

Kidonakis and Sterman '97; R. B., Catani, Mangano, and Nason '98-'03.

- Next-to-Next-to-Leading-Logs (NNLL)

Moch and Uwer '08; Beneke et al. '09-'10; Czakon et al. '09; Kidonakis '09; Ahrens et al. '10

NLO+NLL Theoretical Prediction



TEVATRON

$$\sigma_{t\bar{t}}^{\text{NLO+NLL}}(\text{Tev}, m_t = 171 \text{ GeV}, \text{CTEQ6.5}) = 7.61 \begin{array}{c} +0.30(3.9\%) \\ -0.53(6.9\%) \end{array} (\text{scales}) \begin{array}{c} +0.53(7\%) \\ -0.36(4.8\%) \end{array} (\text{PDFs}) \text{ pb}$$

LHC

$$\sigma_{t\bar{t}}^{\text{NLO+NLL}}(\text{LHC}, m_t = 171 \text{ GeV}, \text{CTEQ6.5}) = 908 \begin{array}{c} +82(9.0\%) \\ -85(9.3\%) \end{array} (\text{scales}) \begin{array}{c} +30(3.3\%) \\ -29(3.2\%) \end{array} (\text{PDFs}) \text{ pb}$$

M. Cacciari, S. Frixione, M. Mangano, P. Nason, and G. Ridolfi, JHEP 0809:127,2008



S. Moch and P. Uwer, Phys. Rev. D **78** (2008) 034003

Measurement Requirements for $\sigma_{t\bar{t}}$

Measurement Requirements for $\sigma_{t\bar{t}}$

Experimental requirements for $\sigma_{t\bar{t}}$:

- **Tevatron** $\Delta\sigma/\sigma \sim 12\% \Rightarrow \sim \text{ok!}$
- **LHC** (14 TeV, high luminosity) $\Delta\sigma/\sigma \sim 5\% \ll \text{current theoretical prediction!!}$

Measurement Requirements for $\sigma_{t\bar{t}}$

Experimental requirements for $\sigma_{t\bar{t}}$:

- **Tevatron** $\Delta\sigma/\sigma \sim 12\% \Rightarrow \sim \text{ok!}$
- **LHC** (14 TeV, high luminosity) $\Delta\sigma/\sigma \sim 5\% \ll \text{current theoretical prediction!!}$

Different groups presented approximated higher-order results for $\sigma_{t\bar{t}}$

- Including **scale dep at NNLO**, NNLL soft-gluon contributions, **Coulomb corrections**

$$\sigma_{t\bar{t}}^{\text{NNLOappr}}(\text{Tev}, m_t = 173 \text{ GeV}, \text{MSTW2008}) = 7.04^{+0.24}_{-0.36} (\text{scales})^{+0.14}_{-0.14} (\text{PDFs}) \text{ pb}$$

$$\sigma_{t\bar{t}}^{\text{NNLOappr}}(\text{LHC}, m_t = 173 \text{ GeV}, \text{MSTW2008}) = 887^{+9}_{-33} (\text{scales})^{+15}_{-15} (\text{PDFs}) \text{ pb}$$

Kidonakis and Vogt '08; Moch and Uwer '08; Langenfeld, Moch, and Uwer '09

- Integration of the Invariant mass distribution at **NLO+NNLL**

$$\sigma_{t\bar{t}}^{\text{NLO+NNLL}}(\text{Tev}, m_t = 173.1 \text{ GeV}, \text{MSTW2008}) = 6.48^{+0.17}_{-0.21} (\text{scales})^{+0.32}_{-0.25} (\text{PDFs}) \text{ pb}$$

$$\sigma_{t\bar{t}}^{\text{NLO+NNLL}}(\text{LHC}, m_t = 173.1 \text{ GeV}, \text{MSTW2008}) = 813^{+50}_{-36} (\text{scales})^{+30}_{-35} (\text{PDFs}) \text{ pb}$$

V. Ahrens, A. Ferroglia, M. Neubert, B. D. Pecjak, L. L. Yang, arXiv:1006.4682

Next-to-Next-to-Leading Order

Next-to-Next-to-Leading Order

The NNLO calculation of the top-quark pair hadro-production requires several ingredients:

Next-to-Next-to-Leading Order

The NNLO calculation of the top-quark pair hadro-production requires several ingredients:

- Virtual Corrections
 - two-loop matrix elements for $q\bar{q} \rightarrow t\bar{t}$ and $gg \rightarrow t\bar{t}$
 - interference of one-loop diagrams

Körner et al. '05-'08; Anastasiou and Aybat '08

Next-to-Next-to-Leading Order

The NNLO calculation of the top-quark pair hadro-production requires several ingredients:

● Virtual Corrections

- two-loop matrix elements for $q\bar{q} \rightarrow t\bar{t}$ and $gg \rightarrow t\bar{t}$
- interference of one-loop diagrams

Körner et al. '05-'08; Anastasiou and Aybat '08

● Real Corrections

- one-loop matrix elements for the hadronic production of $t\bar{t} + 1$ parton
- tree-level matrix elements for the hadronic production of $t\bar{t} + 2$ partons


Dittmaier, Uwer and Weinzierl '07-'08

Next-to-Next-to-Leading Order





$$p\bar{p} \rightarrow t\bar{t} + 1 \text{ jet}$$

- Important for a deeper understanding of the $t\bar{t}$ prod (possible structure of the top-quark)
- Important for the charge asymmetry at Tevatron
- Technically complex involving multi-leg NLO diagrams



$$\sigma_{t\bar{t}+j} \text{ (LHC)} = 376.2^{+17}_{-48} \text{ pb (with } p_{T,jet,cut} = 50 \text{ GeV)}$$

confirmed by G. Bevilacqua, M. Czakon, C.G. Papadopoulos, M. Worek, Phys. Rev. Lett. **104** (2010) 162002



$t\bar{t} + 2j$

S. Dittmaier, P. Uwer and S. Weinzierl,
Eur. Phys. J. C **59** (2009) 625

Next-to-Next-to-Leading Order

The NNLO calculation of the top-quark pair hadro-production requires several ingredients:

● Virtual Corrections

- two-loop matrix elements for $q\bar{q} \rightarrow t\bar{t}$ and $gg \rightarrow t\bar{t}$
- interference of one-loop diagrams

Körner et al. '05-'08; Anastasiou and Aybat '08

● Real Corrections

- one-loop matrix elements for the hadronic production of $t\bar{t} + 1$ parton
- tree-level matrix elements for the hadronic production of $t\bar{t} + 2$ partons

Dittmaier, Uwer and Weinzierl '07-'08

● Subtraction Terms

- Both matrix elements known for $t\bar{t} + j$ calculation, BUT subtraction up to 1 unresolved parton, while in a complete NNLO computation of $\sigma_{t\bar{t}}$ we need subtraction terms with up to 2 unresolved partons.

Next-to-Next-to-Leading Order

The NNLO calculation of the top-quark pair hadro-production requires several ingredients:

● Virtual Corrections

- two-loop matrix elements for $q\bar{q} \rightarrow t\bar{t}$ and $gg \rightarrow t\bar{t}$
- interference of one-loop diagrams

Körner et al. '05-'08; Anastasiou and Aybat '08

● Real Corrections

- one-loop matrix elements for the hadronic production of $t\bar{t} + 1$ parton
- tree-level matrix elements for the hadronic production of $t\bar{t} + 2$ partons

Dittmaier, Uwer and Weinzierl '07-'08

● Subtraction Terms

- Both matrix elements known for $t\bar{t} + j$ calculation, BUT subtraction up to 1 unresolved parton, while in a complete NNLO computation of $\sigma_{t\bar{t}}$ we need subtraction terms with up to 2 unresolved partons.

⇒ Need an extension of the subtraction methods at the NNLO.

Gehrmann-De Ridder, Ritzmann '09, Daleo et al. '09,
Boughezal et al. '10, Glover, Pires '10, Czakon '10

Next-to-Next-to-Leading Order

The NNLO calculation of the top-quark pair hadro-production requires several ingredients:

● Virtual Corrections

- two-loop matrix elements for $q\bar{q} \rightarrow t\bar{t}$ and $gg \rightarrow t\bar{t}$
- interference of one-loop diagrams

Körner et al. '05-'08; Anastasiou and Aybat '08

● Real Corrections

- one-loop matrix elements for the hadronic production of $t\bar{t} + 1$ parton
- tree-level matrix elements for the hadronic production of $t\bar{t} + 2$ partons

Dittmaier, Uwer and Weinzierl '07-'08

● Subtraction Terms

- Both matrix elements known for $t\bar{t} + j$ calculation, BUT subtraction up to 1 unresolved parton, while in a complete NNLO computation of $\sigma_{t\bar{t}}$ we need subtraction terms with up to 2 unresolved partons.

⇒ Need an extension of the subtraction methods at the NNLO.

Gehrmann-De Ridder, Ritzmann '09, Daleo et al. '09,
Boughezal et al. '10, Glover, Pires '10, Czakon '10

Two-Loop Corrections to $q\bar{q} \rightarrow t\bar{t}$

Two-Loop Corrections to $q\bar{q} \rightarrow t\bar{t}$

$$|\mathcal{M}|^2(s, t, m, \varepsilon) = \frac{4\pi^2\alpha_s^2}{N_c} \left[\mathcal{A}_0 + \left(\frac{\alpha_s}{\pi}\right) \mathcal{A}_1 + \left(\frac{\alpha_s}{\pi}\right)^2 \mathcal{A}_2 + \mathcal{O}(\alpha_s^3) \right]$$

$$\mathcal{A}_2 = \mathcal{A}_2^{(2\times 0)} + \mathcal{A}_2^{(1\times 1)}$$

$$\begin{aligned} \mathcal{A}_2^{(2\times 0)} = & N_c C_F \left[N_c^2 A + B + \frac{C}{N_c^2} + N_l \left(N_c D_l + \frac{E_l}{N_c} \right) \right. \\ & \left. + N_h \left(N_c D_h + \frac{E_h}{N_c} \right) + N_l^2 F_l + N_l N_h F_{lh} + N_h^2 F_h \right] \end{aligned}$$

218 two-loop diagrams contribute to the 10 different color coefficients

Two-Loop Corrections to $q\bar{q} \rightarrow t\bar{t}$

$$|\mathcal{M}|^2(s, t, m, \varepsilon) = \frac{4\pi^2 \alpha_s^2}{N_c} \left[\mathcal{A}_0 + \left(\frac{\alpha_s}{\pi} \right) \mathcal{A}_1 + \left(\frac{\alpha_s}{\pi} \right)^2 \mathcal{A}_2 + \mathcal{O}(\alpha_s^3) \right]$$

$$\mathcal{A}_2 = \mathcal{A}_2^{(2 \times 0)} + \mathcal{A}_2^{(1 \times 1)}$$

$$\begin{aligned} \mathcal{A}_2^{(2 \times 0)} = & N_c C_F \left[N_c^2 A + B + \frac{C}{N_c^2} + N_l \left(N_c D_l + \frac{E_l}{N_c} \right) \right. \\ & \left. + N_h \left(N_c D_h + \frac{E_h}{N_c} \right) + N_l^2 F_l + N_l N_h F_{lh} + N_h^2 F_h \right] \end{aligned}$$

218 two-loop diagrams contribute to the 10 different color coefficients

- The whole $\mathcal{A}_2^{(2 \times 0)}$ is known numerically

Czakon '08.

Two-Loop Corrections to $q\bar{q} \rightarrow t\bar{t}$

$$|\mathcal{M}|^2(s, t, m, \varepsilon) = \frac{4\pi^2 \alpha_s^2}{N_c} \left[\mathcal{A}_0 + \left(\frac{\alpha_s}{\pi} \right) \mathcal{A}_1 + \left(\frac{\alpha_s}{\pi} \right)^2 \mathcal{A}_2 + \mathcal{O}(\alpha_s^3) \right]$$

$$\mathcal{A}_2 = \mathcal{A}_2^{(2 \times 0)} + \mathcal{A}_2^{(1 \times 1)}$$

$$\begin{aligned} \mathcal{A}_2^{(2 \times 0)} = & N_c C_F \left[N_c^2 A + B + \frac{C}{N_c^2} + N_l \left(N_c D_l + \frac{E_l}{N_c} \right) \right. \\ & \left. + N_h \left(N_c D_h + \frac{E_h}{N_c} \right) + N_l^2 F_l + N_l N_h F_{lh} + N_h^2 F_h \right] \end{aligned}$$

218 two-loop diagrams contribute to the 10 different color coefficients

- The whole $\mathcal{A}_2^{(2 \times 0)}$ is known numerically

Czakon '08.

- The coefficients D_i , E_i , F_i , and A are known analytically (agreement with known res)

R. B., Ferroglia, Gehrmann, Maitre, and Studerus '08-'09

Two-Loop Corrections to $q\bar{q} \rightarrow t\bar{t}$

$$|\mathcal{M}|^2(s, t, m, \varepsilon) = \frac{4\pi^2 \alpha_s^2}{N_c} \left[\mathcal{A}_0 + \left(\frac{\alpha_s}{\pi} \right) \mathcal{A}_1 + \left(\frac{\alpha_s}{\pi} \right)^2 \mathcal{A}_2 + \mathcal{O}(\alpha_s^3) \right]$$

$$\mathcal{A}_2 = \mathcal{A}_2^{(2 \times 0)} + \mathcal{A}_2^{(1 \times 1)}$$

$$\begin{aligned} \mathcal{A}_2^{(2 \times 0)} = & N_c C_F \left[N_c^2 A + B + \frac{C}{N_c^2} + N_l \left(N_c D_l + \frac{E_l}{N_c} \right) \right. \\ & \left. + N_h \left(N_c D_h + \frac{E_h}{N_c} \right) + N_l^2 F_l + N_l N_h F_{lh} + N_h^2 F_h \right] \end{aligned}$$

218 two-loop diagrams contribute to the 10 different color coefficients

- The whole $\mathcal{A}_2^{(2 \times 0)}$ is known numerically

Czakon '08.

- The coefficients D_i , E_i , F_i , and A are known analytically (agreement with known res)


R. B., Ferroglia, Gehrmann, Maitre, and Studerus '08-'09

- The poles of $\mathcal{A}_2^{(2 \times 0)}$ (and therefore of B and C) are known analytically

Ferroglia, Neubert, Pecjak, and Li Yang '09

Two-Loop Corrections to $q\bar{q} \rightarrow t\bar{t}$

- D_i, E_i, F_i come from the corrections involving a closed (light or heavy) fermionic loop:




- A the leading-color coefficient, comes from the planar diagrams:



- The calculation is carried out analytically using:
 - **Laporta Algorithm** for the reduction of the dimensionally-regularized scalar integrals (in terms of which we express the $|\mathcal{M}|^2$) to the Master Integrals (MIs)
 - **Differential Equations Method** for the analytic solution of the MIs


Laporta Algorithm and Diff. Equations



PUBLIC PROGRAMS

- AIR – Maple package
(C. Anastasiou and A. Lazopoulos, JHEP 0407 (2004) 046)
- FIRE – Mathematica package (A. V. Smirnov, JHEP 0810 (2008) 107)
- REDUZE – C++/GiNaC package
(C. Studerus, Comput. Phys. Commun. 181 (2010) 1293)


Master Integrals for N_l and N_h



18 irreducible two-loop topologies (26 MIs)


R. B., A. Ferroglia, T. Gehrmann, D. Maitre, and C. Studerus, JHEP **0807** (2008) 129.

Master Integrals for the Leading Color Coeff



For the leading color coefficient there are 9 additional irreducible topologies (19 MIs)

Example



$$= \frac{1}{m^6} \sum_{i=-4}^{-1} A_i \epsilon^i + \mathcal{O}(\epsilon^0)$$


$$A_{-4} = \frac{x^2}{24(1-x)^4(1+y)},$$

$$A_{-3} = \frac{x^2}{96(1-x)^4(1+y)} \left[-10G(-1; y) + 3G(0; x) - 6G(1; x) \right],$$

$$A_{-2} = \frac{x^2}{48(1-x)^4(1+y)} \left[-5\zeta(2) - 6G(-1; y)G(0; x) + 12G(-1; y)G(1; x) + 8G(-1, -1; y) \right],$$

$$\begin{aligned} A_{-1} = & \frac{x^2}{48(1-x)^4(1+y)} \left[-13\zeta(3) + 38\zeta(2)G(-1; y) + 9\zeta(2)G(0; x) + 6\zeta(2)G(1; x) - 24\zeta(2)G(-1/y; x) \right. \\ & + 24G(0; x)G(-1, -1; y) - 24G(1; x)G(-1, -1; y) - 12G(-1/y; x)G(-1, -1; y) \\ & - 12G(-y; x)G(-1, -1; y) - 6G(0; x)G(0, -1; y) + 6G(-1/y; x)G(0, -1; y) + 6G(-y; x)G(0, -1; y) \\ & + 12G(-1; y)G(1, 0; x) - 24G(-1; y)G(1, 1; x) - 6G(-1; y)G(-1/y, 0; x) + 12G(-1; y)G(-1/y, 1; x) \\ & - 6G(-1; y)G(-y, 0; x) + 12G(-1; y)G(-y, 1; x) + 16G(-1, -1, -1; y) - 12G(-1, 0, -1; y) \\ & - 12G(0, -1, -1; y) + 6G(0, 0, -1; y) + 6G(1, 0, 0; x) - 12G(1, 0, 1; x) - 12G(1, 1, 0; x) + 24G(1, 1, 1; x) \\ & - 6G(-1/y, 0, 0; x) + 12G(-1/y, 0, 1; x) + 6G(-1/y, 1, 0; x) - 12G(-1/y, 1, 1; x) + 6G(-y, 1, 0; x) \\ & \left. - 12G(-y, 1, 1; x) \right] \end{aligned}$$

Example



$$= \frac{1}{m^6} \sum_{i=-4}^{-1} A_i \epsilon^i + \mathcal{O}(\epsilon^0)$$

$$A_{-4} = \frac{x^2}{24(1-x)^4(1+y)},$$

$$A_{-3} = \frac{x^2}{96(1-x)^4(1+y)} \left[-10G(-1; y) + 3G(0; x) - 6G(-1, -1; y) \right]$$

$$A_{-2} = \frac{x^2}{48(1-x)^4(1+y)} \left[-5\zeta(2) - 6G(-1; y)G(0; x) + 12G(-1, -1; y)G(0; x) \right]$$

$$A_{-1} = \frac{x^2}{48(1-x)^4(1+y)} \left[-13\zeta(3) + 38\zeta(2)G(-1; y) + 9\zeta(2)G(0; x) + \frac{6\zeta(2)G(1; x)}{\rho \xrightarrow{\frac{\rho}{s}} 1} - 24\zeta(2)G(-1/y; x) \right. \\ + 24G(0; x)G(-1, -1; y) - 24G(1; x)G(-1, -1; y) - 12G(-1/y; x)G(-1, -1; y) \\ - 12G(-y; x)G(-1, -1; y) - 6G(0; x)G(0, -1; y) + 6G(-1/y; x)G(0, -1; y) + 6G(-y; x)G(0, -1; y) \\ + 12G(-1; y)G(1, 0; x) - 24G(-1; y)G(1, 1; x) - 6G(-1; y)G(-1/y, 0; x) + 12G(-1; y)G(-1/y, 1; x) \\ - 6G(-1; y)G(-y, 0; x) - 12G(-1; y)G(-y, 1; x) + 16G(-1, -1, -1; y) - 12G(-1, 0, -1; y) \\ - 12G(0, -1, -1; y) + 6G(0, 0, -1; y) + 6G(1, 0, 0; x) - 12G(1, 0, 1; x) - 12G(1, 1, 0; x) + 24G(1, 1, 1; x) \\ - 6G(-1/y, 0, 0; x) + 12G(-1/y, 0, 1; x) + 6G(-1/y, 1, 0; x) - 12G(-1/y, 1, 1; x) + 6G(-y, 1, 0; x) \\ \left. - 12G(-y, 1, 1; x) \right]$$

1- and 2-dim GHPLs

GHPLs

- One- and two-dimensional Generalized Harmonic Polylogarithms (GHPLs) are defined as repeated integrations over set of basic functions. In the case at hand

$$f_w(x) = \frac{1}{x-w}, \quad \text{with } w \in \left\{0, 1, -1, -y, -\frac{1}{y}, \frac{1}{2} \pm \frac{i\sqrt{3}}{2}\right\}$$
$$f_w(y) = \frac{1}{y-w}, \quad \text{with } w \in \left\{0, 1, -1, -x, -\frac{1}{x}, 1 - \frac{1}{x} - x\right\}$$

- The weight-one GHPLs are defined as

$$G(0; x) = \ln x, \quad G(w; x) = \int_0^x dt f_w(t)$$

- Higher weight GHPLs are defined by iterated integrations


$$G(\underbrace{0, 0, \dots, 0}_n; x) = \frac{1}{n!} \ln^n x, \quad G(w, \dots; x) = \int_0^x dt f_w(t) G(\dots; t)$$

- Shuffle algebra. Integration by parts identities

Remiddi and Vermaseren '99, Gehrmann and Remiddi '01-'02, Aglietti and R. B. '03, Volla and Weinzierl '04, R. B., A. Ferroglia, T. Gehrmann, and C. Studerus '09


Coefficient A

Finite part of A



$$\eta = \frac{s}{4m^2} - 1, \quad \phi = -\frac{t - m^2}{s}$$

Threshold expansion versus exact result



$$\beta = \sqrt{1 - \frac{4m^2}{s}}$$

partonic c.m. scattering angle = $\frac{\pi}{2}$

Numerical evaluation of the GHPLs with GiNaC C++ routines.

Vollinga and Weinzierl '04

Two-Loop Corrections to $gg \rightarrow t\bar{t}$

Two-Loop Corrections to $gg \rightarrow t\bar{t}$

$$|\mathcal{M}|^2(s, t, m, \varepsilon) = \frac{4\pi^2 \alpha_s^2}{N_c} \left[\mathcal{A}_0 + \left(\frac{\alpha_s}{\pi} \right) \mathcal{A}_1 + \left(\frac{\alpha_s}{\pi} \right)^2 \mathcal{A}_2 + \mathcal{O}(\alpha_s^3) \right]$$

$$\mathcal{A}_2 = \mathcal{A}_2^{(2 \times 0)} + \mathcal{A}_2^{(1 \times 1)}$$

$$\begin{aligned} \mathcal{A}_2^{(2 \times 0)} = & (N_c^2 - 1) \left(N_c^3 A + N_c B + \frac{1}{N_c} C + \frac{1}{N_c^3} D + N_c^2 N_l E_l + N_c^2 N_h E_h \right. \\ & + N_l F_l + N_h F_h + \frac{N_l}{N_c^2} G_l + \frac{N_h}{N_c^2} G_h + N_c N_l^2 H_l + N_c N_h^2 H_h \\ & \left. + N_c N_l N_h H_{lh} + \frac{N_l^2}{N_c} I_l + \frac{N_h^2}{N_c} I_h + \frac{N_l N_h}{N_c} I_{lh} \right) \end{aligned}$$

789 two-loop diagrams contribute to **16** different color coefficients

- No numeric result for $\mathcal{A}_2^{(2 \times 0)}$ yet
- The poles of $\mathcal{A}_2^{(2 \times 0)}$ are known analytically

Ferrogia, Neubert, Pecjak, and Li Yang '09

- The coefficients $A, E_l - I_l$ can be evaluated analytically as for the $q\bar{q}$ channel

R. B., Ferrogia, Gehrmann, von Manteuffel and Studerus, in preparation

Two-Loop Corrections to $gg \rightarrow t\bar{t}$

$$|\mathcal{M}|^2(s, t, m, \varepsilon) = \frac{4\pi^2 \alpha_s^2}{N_c} \left[\mathcal{A}_0 + \left(\frac{\alpha_s}{\pi}\right) \mathcal{A}_1 + \left(\frac{\alpha_s}{\pi}\right)^2 \mathcal{A}_2 + \mathcal{O}(\alpha_s^3) \right]$$

$$\mathcal{A}_2 = \mathcal{A}_2^{(2 \times 0)} + \mathcal{A}_2^{(1 \times 1)}$$

$$\begin{aligned} \mathcal{A}_2^{(2 \times 0)} = & (N_c^2 - 1) \left(N_c^3 \mathbf{A} + N_c \mathbf{B} + \frac{1}{N_c} \mathbf{C} + \frac{1}{N_c^3} \mathbf{D} + N_c^2 N_l \mathbf{E}_l + N_c^2 N_h \mathbf{E}_h \right. \\ & + N_l \mathbf{F}_l + N_h \mathbf{F}_h + \frac{N_l}{N_c^2} \mathbf{G}_l + \frac{N_h}{N_c^2} \mathbf{G}_h + N_c N_l^2 \mathbf{H}_l + N_c N_h^2 \mathbf{H}_h \\ & \left. + N_c N_l N_h \mathbf{H}_{lh} + \frac{N_l^3}{N_c} \mathbf{I}_l + \frac{N_h^3}{N_c} \mathbf{I}_h \right) \end{aligned}$$

- We finished the evaluation
of the leading-color coefficient
NO additional MI

789 two-loop diagrams contribute to **16** different channels

- No numeric result for $\mathcal{A}_2^{(2 \times 0)}$ yet
- The poles of $\mathcal{A}_2^{(2 \times 0)}$ are known analytically

Ferrogia, Neubert, Pecjak, and Li Yang '09

- The coefficients $A, E_l - I_l$ can be evaluated analytically as for the $q\bar{q}$ channel

R. B., Ferrogia, Gehrmann, von Manteuffel and Studerus, in preparation

Two-Loop Corrections to $gg \rightarrow t\bar{t}$

$$|\mathcal{M}|^2(s, t, m, \varepsilon) = \frac{4\pi^2 \alpha_s^2}{N_c} \left[\mathcal{A}_0 + \left(\frac{\alpha_s}{\pi} \right) \mathcal{A}_1 + \left(\frac{\alpha_s}{\pi} \right)^2 \mathcal{A}_2 + \mathcal{O}(\alpha_s^3) \right]$$

$$\mathcal{A}_2 = \mathcal{A}_2^{(2 \times 0)} + \mathcal{A}_2^{(1 \times 1)}$$

$$\begin{aligned} \mathcal{A}_2^{(2 \times 0)} = & (N_c^2 - 1) \left(N_c^3 A + N_c B + \frac{1}{N_c} C + \frac{1}{N_c^3} D + N_c^2 E_l + N_c^2 N_h E_h \right. \\ & + F_l + N_h F_h + \frac{N_l}{N_c^2} G_l + \frac{N_h}{N_c^2} G_h + N_c N_l^2 H_l + N_c N_h^2 H_h \\ & \left. + N_c N_l N_h H_{lh} + \frac{N_c^2}{N_c} I_l + \frac{N_h^2}{N_c} I_h + \frac{N_l N_h}{N_c} I_{lh} \right) \end{aligned}$$

- For the light-fermion contrib
one missing topology to reduce

- up to now

7 additional MIs

different color coefficients

analytically

Ferrogia, Neubert, Pecjak, and Li Yang '09

The coefficients $A, E_l - I_l$ can be evaluated analytically as for the $q\bar{q}$ channel

R. B., Ferrogia, Gehrmann, von Manteuffel and Studerus, in preparation

Conclusions

- In the last 15 years, Tevatron explored top-quark properties reaching a remarkable experimental accuracy. The top mass could be measured with $\Delta m_t/m_t = 0.75\%$ and the production cross section with $\Delta\sigma_{t\bar{t}}/\sigma_{t\bar{t}} = 9\%$. Other observables could be measured only with bigger errors.
- At the LHC the situation will further improve. The production cross section of $t\bar{t}$ pairs is expected to reach the accuracy of $\Delta\sigma_{t\bar{t}}/\sigma_{t\bar{t}} = 5\%!!$
- This experimental precision requires a complete NNLO theoretical analysis.
- In this talk I briefly reviewed the analytic evaluation of the two-loop matrix elements.
 - The corrections involving a fermionic loop (light or heavy) in the $q\bar{q}$ channel are completed, together with the leading color coefficient.
 - Analogous corrections in the gg channel can be calculated with the same technique and are at the moment under study.
- The calculation of the crossed diagrams and of the diagrams with a heavy loop have still to be afforded.